


BIOSECURITY HANDBOOK

September 2025



Government of South Georgia & the South Sandwich Islands

Contents

Biosecurity Handbook	4
Summary	4
Who is affected by the Processes:	4
1. Introduction	5
What's the risk?.....	6
2. Visitors	8
2.1. General measures.....	9
2.2. Packing guidelines for personal baggage.....	9
2.3. Pre-border biosecurity checks.....	10
2.4. Going ashore	10
2.5. Checks between sites.....	11
2.6. Additional measures for high-biosecurity- risk groups.....	12
2.7. The Biosecurity Audit System	12
3. Vessels.....	14
3.1. General Measures for the Prevention of Rodents	14
3.2 Use of jetties.....	15
3.3 Pharos SG.....	16
3.4 British Antarctic Survey and other UK Science vessels	17
3.5 Royal Navy vessels.....	17
3.6 Yachts	17
3.7 General measures for invertebrate detection and deterrence	18
4. Cargo	19
4.1 Guidelines for packing facilities	19
4.2 Cargo packing requirements.....	19
4.3 Procedure for packing containers	20
4.4 Procedure for cleaning vehicles	20
4.5 Pre-border checks for cargo.....	21
4.6 Border inspections for cargo and personal equipment	22
4.7 Mail and Post	23
4.8 Aggregate	24
4.9 Movement of cargo between King Edward Point and Bird Island.....	24
5. Fresh Produce.....	25
5.1 General measures.....	25
5.2 Guidelines for ordering fresh produce.....	25

5.3 Biosecurity checks on arrival on station.....	26
6. Post-Border Monitoring.....	28
6.1 King Edward Point and Grytviken rodent monitoring procedures.....	28
6.2 Sticky traps for monitoring invertebrate and mouse incursions.....	29
7. Reporting	30
7.1 General Principle	30
7.2 Reporting procedures for visitors.....	30
7.3 Reporting procedures for Government Officers.....	31
7.4 Annual biosecurity review	31
8. Incursion Response.....	32
8.1 Rodent incursion	32
8.2 Non-native plant incursion.....	32
8.3 Non-native invertebrate introduction.....	33
8.4 Marine non-native species	33
9. Highly Pathogenic Avian Influenza	34
9. 1. Background	34
9. 2. Key principals	34
9.3 Activity specific response measures	35
9.4. HPAI Biosecurity	37
9. 5. Signs and symptoms.....	37
9.6. Personal Protective Equipment (PPE) for science activities.....	38
9.7. Bird strike.....	39
9.8. Removal of carcasses	39
9.9 Report form for suspected disease outbreak in wild birds	40
Annex 1 - vehicle cleaning checklist	41
Annex 2 Government of South Georgia & the South Sandwich Islands Number and placement of rodent bait boxes on vessels.....	42

Biosecurity Handbook

Summary

This handbook is designed to provide reference for safeguarding South Georgia & the South Sandwich Islands against the introduction and spread of invasive non-native species and pathogens.

This document draws together all GSGSSI's current biosecurity policies into a single place so they can be easily accessed.

It is widely accepted that one of the greatest threats to biodiversity on island ecosystems is the introduction of invasive non-native species (INNS). With increasing numbers of people visiting South Georgia & the South Sandwich Islands, the risk of introduction of INNS is increased.

In recent years, several large habitat restoration projects have sought to eradicate rodents, reindeer and invasive weeds from South Georgia. Significant resources have been used in these important conservation projects and it is now more important than ever that stringent biosecurity measures are in place to prevent non-native species entering the Territory.

Who is affected by the Processes:

This policy applies to all persons entering and moving within the Territories, any persons involved with the operation of vessels within the Territorial Seas of the Territories, or persons involved with sending cargo to the Territories.

For more information, contact info@gov.gs.

Failure to adhere to this guidance could be used as evidence for any prosecution under the Wildlife and Protected Areas Ordinance (2011 and amendment 2013).

1. Introduction

South Georgia & the South Sandwich Islands are a haven for wildlife and are home to globally important populations of marine mammals and seabirds, including five million seals of four different species, and 65 million breeding birds of 30 different species. Environmental stewardship is embodied within the [Protect, Sustain, Inspire](#) framework which guides the work of the Government of South Georgia & the South Sandwich Islands (GSGSSI).

This Biosecurity Handbook has been developed in response to our strategic commitments, recognising biosecurity infringements represent one of the most significant risks to the future sustainable management of this UK Overseas Territory.

This Biosecurity Handbook represents a statement of the steps that need to be taken to ensure that the Territories remain safe from biosecurity threats. Failure to adhere to this measure outlined in this handbook could be used as evidence for any prosecution under the Wildlife and Protected Areas Ordinance (2011 and amendment 2013).

To make sure this Handbook reflects best practice, GSGSSI undertakes an annual review of biosecurity operations, in conjunction with stakeholders, to identify any emerging threats and take action to mitigate risk including the development of new and improved control measures.

Non-native species can enter South Georgia & the South Sandwich Islands through a range of pathways including ships, in cargo, with passengers and within personal baggage. To mitigate these risks, GSGSSI works across the biosecurity continuum and has procedures in place pre-border, on the border, and post-border to limit the opportunities for non-native species to get to, and establish in the Territories (Fig. 1).




Figure 1. Effective biosecurity operates along the entire biosecurity continuum

The foundation of this process is ensuring that there is effective, comprehensive biosecurity measures in place and that it is accessible to all user groups; the purpose of this Biosecurity Handbook.

What's the risk?

A Horizon Scanning exercise undertaken by the GB Non-Native Species Secretariat, used a consensus approach with the input of over 30 experts in their field to identify a priority species list for South Georgia, based on the likelihood of their arrival, establishment and spread, with each species being subsequently scored for impacts on biodiversity (bio), economy (eco) and human health (hlth).

The following table summarises the species which are likely to be the greatest risks to South Georgia's biosecurity, or to put it another way 'South Georgia's most unwanted'.

Table 1 The greatest threats to South Georgia

Species	Common_names	Group	Impact		
			bio	econ	hlth
<i>Mytilus chilensis</i> (?)	Chilean mussel	Marine	x		
<i>Mytilus edulis</i> (?)	Blue mussel	Marine	x		
<i>Botryllus schlosseri</i>	Colonial Ascidian	Marine	x		
<i>Carcinus maenas</i>	European Shore Crab	Marine	x		
<i>Ciona intestinalis</i>	Ascidian	Marine	x		
<i>Halicarcinus planatus</i>	Decapod	Marine	x		
<i>Mytilus galloprovincialis</i>	Mediterranean mussel	Marine	x		
<i>Undaria pinnatifida</i>	Asian kelp	Marine	x		
<i>Bugula neritina</i>	Ruby bryozoan	Marine	x		
<i>Austromininus modestus</i>	Darwins Barnacle	Marine	x		
<i>Codium fragile</i> subsp <i>fragile</i>	Green sea fingers - Green Alga	Marine	x		
<i>Ascidia aspersa</i>	European sea squirt	Marine	x		
<i>Rattus rattus</i>	black rat	Mammal	x	x	
<i>Mus musculus</i>	house mouse	Mammal	x	x	
<i>Rattus norvegicus</i>	brown rat	Mammal	x	x	
<i>Forficula auricularia</i>	European earwig (from Falklands)	Dermaptera	x		
<i>Hypogastrura manubrialis</i>	springtail	Springtail	x		
<i>Acaena lucida</i>		Plant	x		
<i>Carex trifida</i>		Plant	x		
<i>Leptinella plumosa</i>		Plant	x		

2. Visitors

Masters, Skippers and/or Expedition Leaders must ensure all biosecurity measures described in this Handbook are in place before departing for the Territories. Failure to do so may constitute an offence under the Wildlife and Protected Areas Ordinance (2011) resulting in prosecution and could have catastrophic consequences for the ecology of the Territories.

South Georgia & the South Sandwich Islands has no permanent human population. Everyone who comes to the Territories is therefore a “visitor” and has a responsibility to preserve the environment for future generations. All visitors can help in preventing new non-native species arriving in the Territory or moving existing established non-native species between sites.

Vessels departing Stanley for South Georgia may be asked by the Government to undergo rodent screening by a Biosecurity Detector Dog Team. Typically, a vessel & cargo search takes the dog team less than one hour to perform.

The Government has adopted a risk-based structure for biosecurity dog searches in Stanley. Vessel risk will be assessed by history of searches (residual odours/previous rodents), vessel class, and risk of rodent getting ashore on SG (vessel activities – cruising vs landing vs coming alongside). The GSGSSI Visitor Manager will communicate with the Masters, Skippers and/or Expedition Leaders of the vessel ahead of their call to Stanley to confirm if your vessel will require Biosecurity Detector Dog Team inspection.

Rat guards should be deployed at all times when vessels are alongside at any ports prior to visiting the Territories. Be vigilant to signs of rodents on your vessel before and during your visit and undertake rigorous pre-departure inspections.

Rodent monitoring stations supplied by the Government should be collected from the Government offices in Stanley, Falkland Islands prior to departure for South Georgia. However, if arriving via other ports the vessel may provide their own rodent bait stations conforming with the GSGSSI Rodent bait protocol described in Annex 2 to this Handbook.

Rodent monitoring stations must be checked prior to entering the Territories Maritime Zone, and the findings reported to the Government Officers at King Edward Point. The Government Officer at King Edward Point must be contacted immediately if there is ever any concern or suspicion about the presence of rodents on a vessel. Any vessel believed to have rodents on will need to satisfy GSGSSI that appropriate remedial action has been taken and will be required to leave the Territory until such time as this action has been taken.

2.1. General measures

All persons arriving in the Territories are “visitors”, regardless of the purpose of their visit as a scientist, government personnel, expedition staff, tourist or serving in HM armed forces. It is imperative that all visitors meet biosecurity obligations.

To fully understand the importance of biosecurity, and the measures which should be taken before and during a visit to South Georgia and/or the South Sandwich Islands, all visitors must receive appropriate briefings from their trip organiser and watch the GSGSSI visitor briefing film before arriving in the Territories. Only in extenuating circumstances where this has not been possible, can a Government Officer arrange a verbal briefing and viewing prior to disembarkation. If you believe this may apply to you contact the Government Officers as soon as possible. You will need to head directly for King Edward Point / Grytviken.

2.2. Packing guidelines for personal baggage

Visitors are asked to follow these simple guidelines when packing their personal baggage:

What are you looking for?

- Any foreign organic matter such as soil/dirt/sediment, seeds and plant material, animal material (such as fur, feathers and/or faeces), animal marks including gnaw marks and invertebrates.

Purchasing considerations:

- Where possible, take new clothing and equipment, especially coats, over-trousers, boots and bags.
- Choose outdoor wear without Velcro, webbing/mesh and boots that have open treads which will be easy to clean.
- Avoid woven or knitted gloves and instead opt for smooth waterproof options.
- Plant material can often become caught in the mesh back and waist belt of rucksacks so look for ones with a smooth/tight weave fabric

Before you pack:

- Used clothing and equipment should be washed before leaving home, and any remaining seeds, soil or organic material carefully picked off by hand. Pay particular attention to Velcro, fastenings, seams, folds and pockets.
- Day sacks, camera bags, tripods and walking sticks should be thoroughly cleaned.

Tips:

- A vacuum cleaner will help remove dirt from the inside of pockets, bags, nooks and crannies.
- A needle is useful to pick out seeds stuck in Velcro or in seams.
- A screwdriver is useful to remove soil and seeds from the tread of boots.
- A stiff brush or sticky tape may help to remove seeds from clothing.

After you pack:

- Once packed, bags should be kept closed and stored in a clean area (not in a shed or garage) off the ground. This will reduce the risk of invertebrates or mice crawling inside.

NOTE: Import of fresh produce such as fruit, vegetables, meat, eggs and unpasteurised dairy products for personal consumption onto the islands is not permitted, and no such items should be packed in personal baggage and brought ashore. Poultry products are of particular concern since they may carry avian diseases.

2.3. Pre-border biosecurity checks

Often, visitors travel to a number of other destinations before finally reaching South Georgia and/or the South Sandwich Islands. Therefore, even if the packing guidelines have been followed prior to leaving home, they will need to be repeated before making the first landing in the Territory. It is the responsibility of the Masters, Skippers and/or Expedition Leaders or person in charge of the visit to ensure that all visitors carry out the following actions:

- Thoroughly inspect and clean all luggage and equipment to be brought ashore, such as daypacks and camera bags.
- Special attention should be paid to Velcro, footwear, gaiters, pockets, turn-ups in trousers, and the hoods of jackets (pockets to be turned inside out or vacuumed).
- Daypacks and camera bags must be brushed out and vacuumed to remove soil/dirt/sediment, seeds, and organic material.

2.4. Going ashore

All visitors should note the following when planning a shore excursion:

- Boot washing is obligatory for all persons prior to going ashore and again when returning to the ship. Boots must be cleaned to remove dirt and seeds and then dipped in an approved biocide (e.g. Virkon S or other similar biocides) which is used according to the manufacturers instructions. Masters, Skippers and/or Expedition Leaders must ensure that they have adequate supplies of biocides for boot washing.
- Clothing inspections and boot washing must be overseen by a competent member of the visit team, for tourist visits this must be either a member of the expedition staff or an appropriate crew member. Masters, Skippers and/or Expedition Leaders are responsible for ensuring that this inspection is carried out. All external surfaces of footwear, which will be worn ashore, must be washed.



NOTE: Government Officers will inspect boot washing facilities and procedures on visiting vessels (including yachts) and will carry out inspections of visitors, including staff and crew, before they disembark the vessel to ensure biosecurity protocols have been undertaken properly.

- All boats and tenders must be thoroughly inspected for rodents, invertebrates and organic material before embarking passengers, and again when departing shore to return to the ship.
- No loose cargo should be landed (such as loose items in open bags or nets). All cargo should be inspected, boxed and sealed before landing.
- Bags should not be left open and unattended ashore.
- Visitors should avoid putting day sacks or camera cases down on the ground where they may pick up soil, seeds and invertebrates which could be transferred between sites.
- Fresh fruit, vegetables, meat, eggs and unpasteurised dairy products are not to be taken ashore.
- Any permitted foodstuffs that are brought ashore must be in boxes that are robust and fully sealed. Boxes should be made from and either plastic, metal or wood.

2.5. Checks between sites

Different areas of South Georgia & the South Sandwich Islands are environmentally unique, and it is important not to move material between regions. As well as potentially spreading alien plant or invertebrate species to uninhabited sites, there is the potential to spread disease between colonies of seals and seabirds.

- Boot washing facilities should be cleaned and refilled for each new landing or every time personnel move between regions.
- All personnel must inspect clothing, boots/footwear, luggage and equipment between landings and repeat cleaning procedures to minimize the risk of intra-regional transfer. Boots should be checked after scrubbing and dipping on return to ensure there is no material remaining.

- All personnel are encouraged to check clothing and equipment for invertebrates and rodents after being on board a vessel that is not their usual base.

2.6. Additional measures for high-biosecurity- risk groups

Activities that involve visitors spending a large amount of time ashore such as during science or media projects, overnight trips undertaken by mountaineering expeditions, or by personnel based at King Edward Point, present an increased biosecurity risk (Fig. 4).



Figure 4. Camping equipment presents a particular biosecurity risk

Therefore, the following additional biosecurity procedures should be undertaken:

- Tents should be dry brushed inside and out to remove soil, seeds or invertebrates. If needed, a damp cloth or hose should be used on heavily soiled areas. Pegs should be scraped clean and then dipped in Virkon S or an appropriate biocide.
- Field clothing should be thoroughly cleaned before being used in different areas of South Georgia and/or the South Sandwich Islands. For personnel based at King Edward Point, this should be done in the biosecurity facility.
- Scientific equipment should be thoroughly cleaned according to the manufacturer's instructions. For any equipment that has been in contact with wild birds, mammals, or soil, cleaning protocols must involve a suitable biocide.

For particularly high-risk projects (such as some construction work, expeditions, science/monitoring in sensitive areas), a bespoke biosecurity plan is necessary. If this is required, it will be identified in the permit application process.

2.7. The Biosecurity Audit System


The Biosecurity Audit is a check undertaken by Government Officers, on all vessels in respect of their compliance with biosecurity procedures, or more specifically the effectiveness of the procedures in place to reduce biosecurity risk.

Government Officers conduct a standardised inspection of clothing, footwear and bags to check for biosecurity risks. A tiered system comprising ‘**basic**’ and ‘**comprehensive**’ audits is in place. This system applies to all vessels that disembark crew or passengers at any site in the Territory.

Both types of audit follow the same inspection procedure but for a basic audit the sample sizes and failure thresholds are not intended to be statistically significant, rather they are an early warning sign that a more comprehensive audit is required on the next visit and remedial action should be taken immediately by the vessel. They are scaled by vessel size.

For a comprehensive audit, a statistically significant sample size will be calculated which will be directly proportional to the number of passengers on board. The failure threshold for a comprehensive audit is 95%

At the start of the season, all visiting vessels have a ‘comprehensive’ passenger audit. Providing vessels get above 95% in a comprehensive audit, for subsequent visits, they can have a ‘basic’ audit.



After all audits, feedback if provided to the vessel (EL or responsible person)

If an audit (basic or comprehensive) is failed, on the next visit, a comprehensive audit is undertaken. However, there is no specific requirement to come to KEP/Grytviken first.

All vessels should strive to achieve a 100% pass rate, and many achieve this!

3. Vessels

3.1. General Measures for the Prevention of Rodents

The only way goods and people can access South Georgia & the South Sandwich Islands is by ship. As a result, a wide variety of vessels operate within the Maritime Zone (MZ). The primary risk to terrestrial biodiversity associated with vessels (as opposed to the cargo they carry) is the potential for them to harbour rodents that are subsequently transferred ashore.

The following measures apply to ALL vessels that operate in the SGSSI MZ:

- A valid Ship's Sanitation Certificate must be in date on entry into the Territories.
- Effective rat guards must be fitted to mooring lines when alongside at any port; they should be fitted such that they will not blow off in strong wind, or allow rodents to pass over/around them.
- External doors and windows should be closed whenever possible.
- Rodent bait boxes with non-toxic bait must be carried on board in each of the areas listed in the table below where present. This means that a vessel could require up to 6 bait boxes if each location is present and distinct.

Location	Type	Bait box
Foc'sle (mooring line locker or Bosun's locker)	Rodent entry/exit point	Y/N
Aft mooring deck	Rodent entry/exit point	Y/N
Zodiac storage or shelter deck	Rodent entry/exit point	Y/N
Cargo receiving areas	Rodent entry/exit point	Y/N
Dry food & provision stores	Rodent harbourage	Y/N
Waste storage areas	Rodent harbourage	Y/N

- Rodent monitoring stations must be deployed on all vessels according to Annex 2, before entering the Maritime Zone (MZ). These boxes should be checked and the results reported to the Government Officers upon entry of the SGSSI MZ.
- If rodents are discovered on board, the vessel will be required to leave the MZ for remedial action and a new sanitation inspection. GSGSSI must be satisfied that the infestation is cleared before approval to re-enter the MZ is granted.

NOTE: Rodent monitoring stations are provided to all vessels and may be collected in the GSGSSI office in Stanley, prior to departure for South Georgia, on their first call of the season. Vessels which do not enter the MZ via the Falkland Islands must contact GSGSSI before departure to make necessary arrangements.

3.2 Use of jetties

There are a variety of pathways that a rodent may leave a vessel once in South Georgia. One of the most likely is when a vessel is alongside at a jetty. There are only two jetties on South Georgia, and their use is restricted.

Only vessels discharging or loading cargo (or for other, agreed activities or circumstances such as health and safety issues which require them to be alongside) should be alongside the jetties at KEP and Grytviken.

The jetty at King Edward Point may be used by the vessels listed below, when there is an operational, logistics, or safety reason as follows:

- The GSGSSI fisheries patrol vessel *Pharos SG*.
- GSGSSI/BAS harbour launches (permanently based on South Georgia).
- The BAS vessel *RRS Sir David Attenborough*.
- Royal Navy vessels as agreed.
- Yachts (small sailing or motor vessels) with a load line length of 24 m or less at the discretion of the Government Officer.
- In emergency situations where a serious mechanical failure threatens the safety of those on board.
- Vessels with specific permission detailed on a Regulated Activity Permit or with special permission from GSGSSI, and where a biosecurity risk assessment has been conducted.

The Tijuca jetty at Grytviken may only be used by passenger tenders and yachts (small sailing or motor vessels) with a load line length of 24 m or less.

Use of the jetty is allowed for the above vessels as it is deemed operationally necessary to discharge cargo, transfer personnel, conduct vessel maintenance, allow for crew rest periods or vessel safety during inclement weather. The following biosecurity measures must be in place when a vessel is alongside:

- Gangways should be lifted at night where it is possible to do so. Where this is not possible (i.e. because 24h access is needed to the vessel), consideration should be given to lighting the gangway to reduce the chances of it being used by rodents. However, this must be weighed against the chances of a bird strike.

Because of the increased likelihood of transferring rodents when alongside, vessels which are authorised to use jetties are required to have enhanced biosecurity measures in place, which are described in the relevant sections below.

3.3 Pharos SG

The *Pharos SG* is the Government fishery patrol vessel (Fig. 5) and is engaged in a range of tasks including delivery of cargo and personnel to the station at King Edward Point and sometimes Bird Island, undertaking at sea inspections of fishing vessels and supporting a range of environmental, heritage and scientific projects at sites around the island. Because of this diverse range of tasks, and because the vessel is often working close to the shore at a range of locations, it is imperative that the highest standards of biosecurity are in place at all times on the vessel. The following additional measures are in place on board:

- The ship will be subject to a search a maximum of 24 hours prior to departure by the rodent detection dog team in the Falklands.
- The ship's company is regularly briefed on biosecurity policy and any particular biosecurity considerations for upcoming tasking.
- Rodent control stations are placed at FIPASS, the home port in the Falklands, to limit the local abundance of rodents and reduce the likelihood of one boarding the vessel.
- Rodent bait stations are placed at key locations on the vessel which are checked by the Master before arriving in the South Georgia Maritime Zone and reported to the Government Officers at KEP. These should be kept under regular review in conjunction with the GSGSSI rodent biosecurity team.
- Insect traps (sticky and UV) are placed in food storage areas and accommodation areas and are regularly checked by the master. Traps should be replaced at a minimum every 6 months and any found insects of concern (such as cockroaches, earwigs, springtails, etc.) should be reported to the Government Officers. Fumigation may be necessary to treat any infestation.
- When transferring observers or fishery officers between vessels, personal clothing and equipment is required to be checked and cleaned immediately upon arrival on board. If there is any suspicion that the previous vessel had an invertebrate infestation, these checks should be made on deck, not inside the accommodation space. Insecticide should be on hand.



Figure 5. Pharos SG the GSGSSI fisheries patrol vessel

3.4 British Antarctic Survey and other UK Science vessels

British Antarctic Survey vessel *RRS Sir David Attenborough* visits South Georgia during annual resupply operations and in support of science projects. During annual resupply, a large volume of cargo is discharged, and the vessel may be alongside for several days and so enhanced biosecurity procedures are required. These include:

- The ship will be subject to a search a maximum of 24 hours prior to departure by the rodent detection dog team in the Falklands.
- Personnel traveling on the BAS vessels are briefed on environmental matters, including biosecurity, prior to travel. Where necessary this can be arranged through GSGSSI in Stanley.
- At least six rodent bait stations are placed at key locations on the vessel which are checked by the Master on entry into the Maritime Zone, and the day before the vessel comes alongside at KEP and the findings reported to the Government Officers.
- Sticky and UV insect killers are placed inside all food storage areas.
- Ultrasonic rat deterrents are placed on gangways.
- Rat guards must be in place at all gateway ports immediately upon arrival.

3.5 Royal Navy vessels

A range of Royal Navy vessels visit the Territory and may disembark passengers at a range of sites. Biosecurity is important at all sites.

Regardless of whether Royal Navy vessels require the use of the jetty at King Edward Point or not, to mitigate the risk of transferring rodents, the following enhanced biosecurity measures are in place:

- The ship will be subject to a search by the rodent detection dog team a maximum of 24 hours prior to departure from the Falklands.
- Military vessels will appoint a 'Biosecurity point person' who will be responsible for ensuring that the vessel's visit, complies with the biosecurity procedures detailed in this Biosecurity Handbook.
- Rodent control stations are placed at Mare Harbour, the home port in the Falklands, to limit the local abundance of rodents and reduce the likelihood of one boarding the vessel.
- The vessel will maintain best practices at gateway ports, including the use of rodent guards on mooring lines, to reduce the risk of a rodent getting on board.
- Rodent bait stations are placed at a minimum of six key locations on the vessel and must be checked by a designated Officer prior to entry into the South Georgia Maritime Zone and the findings reported to the Government Officers.

3.6 Yachts

A range of yachts visit South Georgia & the South Sandwich Islands each year. .

- Yachts departing from the Falkland Islands for the Territories will be subject to a search by the rodent detection dog team a maximum of 24 hours prior to departure..
- All yachts must have suitable rodent monitoring bait stations. We recommend two for yachts under 24 m. Bespoke advice can be provided upon asking. Yachts travelling to the Territory via the Falklands can collect rodent monitoring stations and receive a briefing from the GSGSSI office in Stanley if required. Yachts not passing through the Falklands must source rodent monitoring stations that meet the requirements set out in Annex 2 prior to entering the South Georgia Maritime Zone.
- Rodent bait stations must be checked before entry into the SGSSI MZ and findings reported to the Government Officer.

3.7 General measures for invertebrate detection and deterrence

Invertebrates such as cockroaches, earwigs, spiders, moths and flies can enter vessels in association with personal baggage, when loading provisions or simply by flying on board when the vessel is in port. The large variety of microhabitats on board ships means that once on board, some species can thrive. To reduce the likelihood of invertebrates entering the vessel initially, and to reduce the risk of subsequent transfer between vessels or to the shore, the following measures should be adopted:

- When in a foreign port, where possible, windows and doors should be closed and deck lighting kept to a minimum.
- Cardboard packaging, especially on fresh produce, should be minimised and incinerated as soon as possible.
- Crawling and flying insect traps should be fitted in high-risk areas such as food storage areas and checked regularly so that infestations can be detected early and dealt with.

Government officials may ask to inspect invertebrate traps when on board. If an infestation is discovered, additional biosecurity measures may be required to prevent spread to other vessels or the shore.

4. Cargo

4.1 Guidelines for packing facilities

Reducing the likelihood of seeds, soil, invertebrates and rodents becoming entrained in cargo at the source destination is a vital step in the biosecurity continuum. If a packing facility is clean and free from potential contamination sources, it is likely that the cargo will also remain clean. For commercial packing facilities that ship materials to the Territory, the following guidelines apply:

- There should be active rodent control in the warehouse facility and surrounding area. We are aware of rodent activity in the warehouse and all protocols have been updated to align with rodent activity. Trapping and monitoring protocols are in force. All sending cargo should ensure that cargo is packed rodent proof.
- Seeds can sometimes be brought into warehouses on equipment or on air currents and then become entrained in cargo. Efforts should be made to reduce the densities of weed species around warehouse facilities. This can most easily be achieved using a suitable herbicide.
- If crawling invertebrates access packing materials, they can lay eggs that are hard to detect and pose a biosecurity threat. Crawling and flying insect traps should be installed in packing facilities.
- Cargo should be stored inside wherever possible. Ideally, it should be stored in a dedicated biosecurity area/room/container.
- If stored outside, cargo should be placed on a hard standing to avoid contamination with soil. Cargo should not be placed under overhanging vegetation or trees such that organic material might fall onto the cargo. If items become contaminated with soil and organic material, they should be cleaned before onward transport.
- Warehouse doors and windows should remain shut as far as possible, but especially at night when flying insects may be attracted to the lights.

4.2 Cargo packing requirements

The type of packing material and storage location can have a significant effect on the biosecurity risk posed by cargo items. To reduce this risk, the following requirements should be met:

- For routine cargo operations on the Pharos SG rodent-proof pallet boxes made from high-density plastic will be utilized as far as possible.
- Plastic or metal boxes should be used if possible and cleaned between each use.
- If cardboard boxes are to be used, they should be in good condition (or new if possible) and sealed using packing tape on ALL edges and across all potential openings to make them impenetrable.
- As far as is practicable, all wood packaging likely to be off-loaded in the Territory (such as cases, crates, dunnage, pallets and timbers for the purpose of bracing, separating, protecting or securing cargo) should be new and comply with the International Standards for Phytosanitary Measures No. 15 (ISPM 15). Wooden

packaging that remains on the ship (i.e. is not offloaded) does not have to comply with ISPM 15.

- No soil, moss, used sacking, hay, straw, chaff, or wood shavings shall be used in packing materials. Acceptable alternatives include shredded paper, vermiculite, bubble wrap, and other air-filled cushioning materials.
- Cargo will, where logistically possible, be subject to a search by the rodent detection dog team prior to departure from the Falklands.

4.3 Procedure for packing containers

Shipping containers are universally used to transport goods around the world and some simple precautions can vastly reduce the biosecurity risk they pose. The following standards should be met for any shipping containers that are being transported directly to the Territory:

- Containers should be in good condition with effective door seals that would prohibit the passage of rodents and invertebrates.
- Containers should be cleaned before being packed. Ideally, this should be done using a commercial steam cleaning service, but otherwise being swept out and an insecticide spray used in the corners and along the door threshold will suffice.
- Container doors should not be left open and unattended at any time.
- Containers should be placed on hard standing to prevent soil and mud, and organic material from contaminating the container and cargo.
- Cargo should be packed so that cargo can't fall against the doors and prevent re-closure in case propagules are discovered on opening.
- Once cargo loading is complete, before the container is sealed for shipping, a rodent bait station and crawling insect trap should be placed inside next to the door so they can be easily accessed on opening.
- Unless it contains food, the container should be fumigated with a pyrethrum-based insecticide fogger prior to being sealed.
- The number of containers with food should be minimised i.e. not spread across multiple containers, to reduce the number of un-fumigated containers that are transported.
- Unless required by Customs Officers, sealed containers should not be opened en route to the Territory.
- Containers will, where logistically possible, be subject to a search by the rodent detection dog team prior to departure from the Falklands.

4.4 Procedure for cleaning vehicles

Biological material and soil can become attached to vehicles such as quad bikes, 4x4 vehicles such as Land Rovers, construction vehicles, bicycles etc. during everyday use. When vehicles are moved from one location to another, these materials may also be transferred. The following procedures should be employed to reduce the risk of biological material being transported to the Territory:

- Vehicles should be inspected to ensure that they are free of visible soil and biological material (e.g. plant fragments, seeds, and insects) and if necessary, thoroughly cleaned before being loaded onto the re-supply ship. This should include all external and internal surfaces as well as the undercarriage.
- Where practicable, high-pressure steam/hot water cleaning of vehicles is recommended. Alternatively, vehicles may be cleaned manually, such as with a bucket of water and a brush.
- Any external surfaces of the vehicle that come into contact with the ground i.e. tyres, tracks, skis should be washed with Virkon S or an appropriate biocide.
- Vehicle interiors, upholstery and mats should be brushed and/or vacuum cleaned to remove any soil or biological material.
- Engine compartments should be carefully checked for the presence of biosecurity risks such as rodents, seeds, invertebrates and organic material.
- The interior of vehicles should be fumigated with a pyrethrum-based insecticide prior to being shipped.
- If being shipped in a container the container should also be fumigated with a pyrethrum-based broad spectrum insecticide.

Any vehicles being transported through the Falkland Islands, should be presented to a GSGSSI official with at least 5-working days before cargo loading in order for checks to be carried out and any necessary remedial measures carried out. Consignees should work through the vehicle cleaning checklist and sign the declaration (see annex 1).

If vehicles are not being transported through the Falkland Islands, the shipper should contact GSGSSI to make alternative arrangements. Note that the Falkland Islands may have their own protocols and advice should be sought from the Biosecurity Officer at the Department of Agriculture(biosecurity@doa.gov.fk) to ensure that these can be met before importing a vehicle.

4.5 Pre-border checks for cargo

The majority of cargo on the British Antarctic Survey vessel and some fully-loaded containers of construction material from SATLAN are consigned directly from the UK to South Georgia. Other cargo is either purchased in the Falkland Islands or arrives in the Falklands from the UK as consolidated cargo and needs to be re-packed for onward transport. Under these circumstances there is the potential for non-native species to become entrained with the cargo whilst in storage in the Falkland Islands so a GSGSSI official must undertake some additional checks prior to loading. These include:

- Making sure packaging meets packing guidelines (see 4.3).
- Checking to ensure packaging is still intact and repairing any holes if needed. The contents of any boxes which have been breached will require inspecting to ensure that rodents or invertebrates have not been able to enter.
- Ensuring the inside of containers is clean and the outside of containers have been washed.

If cargo is received at the dockside and a biosecurity risk is identified, the GSGSSI official has the authority to prevent it from being loaded onto the vessel until remedial action has been undertaken.

4.6 Border inspections for cargo and personal equipment

Although every effort is made to prevent non-native species from being accidentally brought into the Territory, it is important that cargo and personal equipment receives a final inspection on arrival in the Territory. During routine operations i.e. not annual resupply, the following protocols apply:

- Smaller items of cargo from Pharos SG and personal bags must be taken straight to a designated biosecurity facility on arrival. There are dedicated biosecurity facilities at both King Edward Point and Bird Island.
- Post bags are taken to the post office, where crawling insect traps and insecticide spray are on hand. A separate Post Office biosecurity SOP is available for postal officers, included in section 4.7.
- On arrival in the biosecurity facility, bags and boxes are opened and thoroughly checked for soil, organic material, seeds, and invertebrates.
- Individuals must sign to say they have completed these checks before the item is released.
- If an item of cargo contains a small amount of biological material, it may be possible to take remedial action on site to reduce the biosecurity risk. The Government Officer is responsible for making this decision, but will consult with the Biosecurity Officer to determine appropriate mitigation measures.
- If an item is heavily infested, it should be sealed inside a container. The Government Officer will identify the best method for destroying or treating the item in conjunction with the Operations Manager and/or Environment Manager.

At the current time, facilities do not exist to place entire shipping containers inside a biosecurity area. To ensure appropriate biosecurity checks can be undertaken, the following measures should be in place.

- When shipping containers come ashore, they should remain sealed until there is time/space to unpack them fully.
- At no point should a partly unpacked container be left open and unattended.
- A Government official should be present when containers are first opened to monitor for any non-native species and check the rodent and invertebrate traps within.
- In the event that a biosecurity hazard is detected, the container should be sealed immediately and returned to the ship for remedial action or removal from the territory.

Once a year, the British Antarctic Survey vessel *RRS Sir David Attenborough* re-supplies the stations at KEP and Bird Island with the majority of the food and equipment needed for the year. Because of the large quantities of cargo, not everything can be processed within the usual biosecurity facility.

During re-supply the following protocols are in place:

- **Containers** – First overpack/cargo unit out of each container should go to biosecurity shed and be fully unpacked and checked. If no breaches are found, the rest of the container can be released for unpacking in the boatshed. If any breaches detected in the first unit, a further overpack/cargo unit to be inspected. If breaches found in second unit, entire container to be fully checked/unpacked via biosecurity shed.
- **Breakbulk** – GO's to identify 10% of breakbulk cargo to be fully unpacked and checked in the biosecurity shed. If breaches are detected, a further 10% to be checked in the biosecurity shed. If fails are detected at this point, 100% checks are to be initiated.
- **Frozen cargo** - can be transferred directly to the cold store
- **P-boxes, B-bags, and ski equipment** – this should be dealt with as personal equipment and should all be checked in the biosecurity shed.

Once the cargo has been cleared, it can be processed through the boatshed, and additional biosecurity devices are installed for the resupply period. These include:

- At least five sticky invertebrate/mouse traps stationed around the periphery of the building.
- Four additional rodent bait stations positioned within the building

Staff are briefed to remain vigilant for biosecurity breaches, and a box containing insecticide spray is available. If a suspicious box is found, it should be taken to the biosecurity shed before it is completely unpacked, and the magnitude of the infestation is established.

4.7 Mail and Post

- Prior to sending mail to South Georgia, it should be packaged biosecurely, notwithstanding any further directions in the Biosecurity Handbook, including under cargo procedures. This means that all parcels should be packaged free from any organic matter. Where dry food is packaged, this should remain in its original packaging. When parcels or letters are closed, they should be completely sealed using tape. This is so that no insects can crawl in. Any small holes in parcels must also be taped close.
- When mail lands in South Georgia, the mail bag should be opened in a biosecure facility, not the post office. Post Officers can then inspect individual mail to ensure it is still biosecurely packaged (as above).
- All mail that passes the inspection can be moved to the post office.
- All mail that fails the inspection, because it was poorly packaged or got damaged in transit, can be arranged to be collected and should be opened in a biosecure facility by the recipient.

4.8 Aggregate

Aggregate is defined as any coarse particulate material used in construction, including sand, gravel, crushed stone, boulders, pebbles, or slag. It may present a biosecurity risk because biological material such as seeds, soil, and invertebrates can become entrained during production and transport.

Decisions will be made on a case-by-case basis, following an Environmental Impact Assessment which includes consideration of biosecurity, or if it is appropriate to source aggregate locally or if it should be imported from outside the Territory.

4.9 Movement of cargo between King Edward Point and Bird Island

On occasion, goods are shipped between the research stations at King Edward Point and Bird Island. King Edward Point is home to several species of introduced plants and invertebrates that are not found on Bird Island. Bird Island is home to dense aggregations of wildlife, and there is potential for the spread of disease. For any cargo/equipment which has been brought ashore and opened/used on station, or cargo which has been brought ashore and stored outside, extreme caution must be exercised and the following steps taken to reduce risk:

- Prior to loading on a re-supply vessel, cargo should be checked and, where necessary, unpacked, cleaned, and re-packed in clean, intact packaging.
- At King Edward Point and Bird Island, small items should be checked in the biosecurity facility.
- Any items which are too large to fit inside a biosecurity facility must be checked on board the vessel.
- Personal effects should be checked and cleaned following the methods described in section 2.3.

5. Fresh Produce

5.1 General measures

Fresh produce such as vegetables, fruit, salad etc. poses a biosecurity risk as it may contain soil, seeds, pests, or diseases. No fresh produce should be brought ashore at any location other than King Edward Point and Bird Island station.

At King Edward Point, the Government Officers are responsible for ensuring the biosecurity checks are correctly implemented and deciding if an infested shipment should be returned to the supplier or if it can be dealt with locally. At Bird Island, the Station Leader fulfils this role.

5.2 Guidelines for ordering fresh produce

Only quantities of fresh food produce sufficient to meet the station's needs between provisioning calls are to be ordered to reduce the burden of biosecurity checking for station personnel.

It has been deemed that some fresh produce items cannot be checked adequately on receipt in the biosecurity facility and therefore should not be ordered. This includes loose leafy vegetables such as:

- Broccoli
- Cauliflower
- Lettuce
- Kale
- Spinach
- Cabbages (white cabbage and red cabbage are acceptable providing the outer leaves are removed)
- Leeks
- Globe artichokes
- Celery
- Pineapples
- Fresh herbs

NOTE: The list is illustrative, not exhaustive. Check with GSGSSI if in doubt.

Root vegetables should only be ordered if they are pre-washed and do not contain surface soil.

Cruise ships may gift fresh produce; this policy applies, and such donations should not be accepted unless they are pre-washed and do not contain soil. The Government Officer should ensure that cruise ships are aware of this policy and do not send fresh produce ashore without authorisation.

5.3 Biosecurity checks on arrival on station

Upon arrival, all fresh produce must be taken directly to a biosecurity facility where it should be checked for signs of infestation by fungus, non-native invertebrates, soil from outside the territory, or non-native plant seeds.

At King Edward Point and Grytviken, produce should be taken to the biosecurity shed (Fig. 8), which is equipped with crawling insect traps and UV flying insect traps. At Bird Island, produce should be kept in the biosecurity shed, which has been prepared prior to offloading.



Figure 8. Checking fresh produce in the biosecurity shed at King Edward Point

If a shipment is heavily infested, the responsible officer should determine if the produce should be returned to the sender/re-supply ship for disposal/incineration. If only a few items are affected, the responsible officer may decide that remedial action can be taken to reduce the biosecurity risk. Depending on the type of infestation the following actions may be taken:

- If an item has fungal growth, the infected part should be cut out and stored until such time as it can be rendered inert*.
- If live invertebrates are found, they should be killed immediately with insecticide spray or placed in ethanol. Be aware that if live or dead invertebrates are seen, there may also be microscopic eggs that could hatch at a later date (timing will depend on species and temperature). Extra vigilance should be exercised, including weekly checks of sticky traps in food storage areas.
- If there are non-native plant seeds associated with the packaging or outer surface of the produce, they should be rendered inert. Seeds may be stored in a sealed container and batch processed if necessary.
- For items with a small amount of surface soil, it may be removed either by removing the skin/peel or by using a dry brush. The material should be rendered inert*.
- Under no circumstances should food or food scraps be fed to local birds.

Onions and garlic should have all outer skins removed, such that only the edible bulb remains. The discarded skins should be rendered inert*.

A log of all biosecurity checks and biosecurity breaches should be kept. The responsible officer is to take photographs of the consignment for evidence if the fresh produce is to be returned to the supplier.

Once processed in the biosecurity facility, fresh produce should be taken to a designated food storage area, i.e., food store, kitchen.

*At King Edward Point, material which needs to be rendered inert should be sent to the Falkland Islands for incineration. Waste will be kept double bagged in the freezer before being shipped to Stanley. At Bird Island, material which needs to be rendered inert should be heated in a pressure cooker for 10 minutes.

6. Post-Border Monitoring

6.1 King Edward Point and Grytviken rodent monitoring procedures

Although the focus of biosecurity efforts is to prevent non-native species from entering the Territory, the vital last stage in the biosecurity continuum is to monitor the efficacy of those efforts and be in a position to take action in the event of an incursion. In the case of rodents, this is particularly important as early detection and rapid response has a high probability of success.

As the majority of visitors and cargo come ashore at King Edward Cove, and it is the only place ships are allowed to come alongside, stringent biosecurity and monitoring are in place. The following monitoring devices have been installed in this area:

- At KEP, there are a minimum of 17 rodent monitoring stations consisting of a DOC 200 trap, an oil-soaked gnaw stick, a wax tag, and a non-toxic bait block.
- At the KEP jetty, there are a minimum of 3 'rat hotels' (Fig. 9).
- A line of at least 24 wax tags radiating out at 25 m intervals from the jetty through the station complex.
- At Grytviken, there are at least 14 rodent monitoring stations consisting of a DOC 200 trap, an oil-soaked gnaw stick, a wax tag, and a non-toxic bait block.
- Around Grytviken, there are a minimum of 2 'rat hotels' (one close to Tijuca, the others(s) at zodiac landing sites).



Figure 9. Rat hotels have a DOC 200 trap, mouse traps, wax tags, poison bait (formulated for rats and mice), and a non-toxic bait block

The Government Officers are responsible for checking the rodent monitoring devices every 2 weeks provided they are accessible.

The Government Officers keep a record of checks that are made.

6.2 Sticky traps for monitoring invertebrate and mouse incursions

Occasionally, invertebrates arrive in cargo, personal baggage, or mail. Although every effort is made to detect these creatures in the biosecurity shed, their cryptic nature means they can escape into the station facilities. They are often then contained within a building, and because only a single animal is present, they do not reproduce and spread. However, there is the potential for multiple individuals or species that are capable of colonising the natural environment to be introduced. Earwigs are prolific in the Falkland Islands, and it is thought that South Georgia could provide a suitable environment for their breeding success. The most effective traps to detect crawling insects are also effective at trapping mice and so serve a dual purpose (Fig. 10). Sticky traps are placed on the ground, alongside walls in areas where the Government Officers deem the risk of an incursion to be highest:

- Crawling insect traps are installed in all inhabited buildings, food and waste storage areas.
- Particular attention is paid to rooms that receive large amounts of luggage, cargo or mail, including the museum storerooms and workshops where building supplies are stored.
- Traps are checked by the Government Officers each month and recorded.



Figure 10. Sticky traps used to monitor for invertebrates and mice

7. Reporting

7.1 General Principle

Keeping records of the number and type of biosecurity inspections undertaken and any incursions that are detected is vital for accurate reporting. In turn, this forms the basis of regular biosecurity reviews that are necessary to ensure that policies remain fit for purpose (Fig. 11).




Figure 11. Biosecurity cycle

All visitors have a responsibility to undertake biosecurity checks and report findings. Findings are typically reported to the Government Officer who collates this information and reports to the Head of Environment and Operations Manager. A Biosecurity Working Group which will act as the forum for reviewing policy, ensuring that procedures are in place to mitigate biosecurity threats facing the Territories will be held every quarter.

7.2 Reporting procedures for visitors

The first and most important element of reporting rests with individuals themselves. Depending on the nature of the visit and a visitor's role on the island there are different reporting requirements:

- Every individual is expected to adhere to GSGSSI biosecurity procedures. We no longer require the signing of the self-audit checklist.
- Masters, Skippers and/or Expedition Leaders (or equivalent) are responsible for ensuring self-audit checks have been completed by all and pass these on to the Government Officer.
- Ships Masters are responsible for ensuring rodent monitoring stations on their vessel (see section 3) are checked and reporting findings to the Government Officer on arrival into the MZ. For vessels that are in the MZ for an extended stay

(e.g. fishing vessels, yachts, Pharos SG), after the initial report, Masters are also responsible for ensuring that bait stations are checked and findings reported to the Government Officers on the 1st and 15th of each month for the duration on their visit.

- All visitors should report any suspicious sightings or signs of rodent or reindeer, and any invertebrates or plants not already known from the islands to the Government Officers via the Masters, Skippers and/or Expedition Leaders (or equivalent). As much information as possible should be included such as a photograph and GPS location.

7.3 Reporting procedures for Government Officers

Government Officers are responsible for consolidating biosecurity reports/ and the day-to-day monitoring of compliance with biosecurity policies at KEP and Grytviken. As part of this role, they are to keep records of checks made on the various biosecurity devices installed around station and ensure that electronic records are completed accurately.

Breaches involving a rodent or invertebrate incursions would be reported immediately.

Biosecurity related incidents which are reported through the British Antarctic Survey Accident Incident Near Miss Environment (AINME) system should be forwarded to GSGSSI from the BAS Environment Office as appropriate.

7.4 Annual biosecurity review

Biosecurity will be reviewed on an annual basis by the Biosecurity Working Group, which includes all members of GSGSSI who have biosecurity within their portfolio of responsibility. This will involve gathering all records together from all parts of the biosecurity continuum and consolidating these into a single document. As part of the review, items and/or activities that have been the most frequent causes of breaches are to be identified and action plans developed to mitigate this risk in the future. The report contains the following sections:

- Pre-departure biosecurity checks.
- Cargo at King Edward Point.
- Intra-regional movements.
- Rodent Monitoring.
- Insect monitoring.
- Biosecurity Audit results.

The Biosecurity Working Group will work together to ensure that action plans are implemented and, where needed, that biosecurity policy is updated. For major changes to the biosecurity policy, appropriate stakeholder consultation will be held.

8. Incursion Response

8.1 Rodent incursion

The South Georgia Heritage Trust (SGHT) led a successful project to eradicate rodents from South Georgia. In March 2011, bait was spread on the Thatcher, Greene, and Mercer Peninsulas and Teie Point. Systematic monitoring undertaken in the three years since baiting indicate that the operation was a success and it is considered that these areas are now rat free. In 2013 SGHT returned to South Georgia to spread bait over the rat infested areas to the north of the Phase 1 area (from the Busen area to Peggotty Bluff). In addition, areas infested with mice (Cape Rosa and the Nunez Peninsula) were baited in 2013. The remaining rat infested areas (to the south of the Phase 1 area from the Barff Peninsula to Cooper Bay) were baited in 2015.

The SGHT eradication project involved a significant amount of planning and the cost was in the region of £10 million. It is therefore vital that the right protocols are in place to reduce the risk of any future rodent incursion. GSGSSI seeks to continue to develop and enhance these protocols, changes to which will be published in future editions of the Biosecurity Handbook.

Most effort should be focused on ensuring that rodents do not enter the Territory, and this Biosecurity Handbook details the measures GSGSSI has in place to prevent any reintroduction. However, in the event that a rodent does get ashore, a comprehensive incursion response plan has been prepared that details actions to be taken to prevent rodents from becoming established and spreading beyond the point of introduction.

This incursion response would be led by Government Officers who would coordinate the response using personnel available on station. The plan is practiced each year to ensure that all the equipment is in place and staff are familiar with procedures. The plan is available from GSGSSI on request.

8.2 Non-native plant incursion

GSGSSI is engaged in a programme to manage non-native plants in the Territory. With the assistance of UK Government funding through the Darwin initiative, great progress has been made in controlling target species around King Edward Point and Grytviken and limiting the spread of species elsewhere. Recently, GSGSSI committed £250,000 over 5 years to continue this work, and it is vital that this is not undermined by the introduction of new species.

As with other groups, the focus of biosecurity is to ensure that new species are not introduced. With plants, this is particularly important because their cryptic nature means in some cases, it may be some time before they are discovered by which time a seed bank could have established.

With the support of the Darwin funding, GSGSSI has prepared a non-native plant management strategy that, in addition to detailing control measures for existing species,

details the process on discovering a new unidentified plant species and the steps needed to make a decision on how to manage the incursion.

8.3 Non-native invertebrate introduction

Plans for dealing with an incursion of non-native invertebrates are focused on measures which can be taken inside accommodation blocks and storage areas. The response will be tailored to the specific invertebrate species that has been detected but would likely involve fumigation followed by a period of intense trapping.

8.4 Marine non-native species

A Blue Belt funded project is underway to assess the risk of marine non-native species and develop suggestions of how GSGSSI may minimise any potential risk.

9. Highly Pathogenic Avian Influenza

9. 1. Background

Avian influenza is a viral disease that primarily affects birds. Low pathogenic avian influenza viruses are common in wild birds and often cause no signs of disease. However, some strains of the virus including H5 and H7 are highly pathogenic in domestic poultry and can cause high mortality if they escape into wild bird populations. These are known as highly pathogenic avian influenza (HPAI). Further H5 and H7 HPAI can emerge from low pathogenicity versions following replication in some species although the low to high pathogenicity switch is predominantly thought to occur in poultry.

The current outbreak of H5N1 HPAI began in 2022 and has resulted in the deaths of high numbers of seabirds in the Northern Hemisphere, the south of Africa and around the Atlantic and Pacific Oceans and throughout South America. Whilst predominantly still a virus that affects birds, during the current outbreak, some mammals have been infected. Often this is due to predators and scavengers consuming infected birds or carcasses, but cases have also been noted in some marine mammals where this mode of transmission would not apply. The hypotheses for the infection of such marine mammals includes environmental exposure or behaviours that may enable exposure within the aquatic environment (e.g., 'playing' with infected carcasses).

In October 2023, HPAI was confirmed in brown skua on Bird Island, South Georgia and since then has been detected at numerous other sites around the Territory and in a range of species including elephant seals, fur seals, wandering albatross, gentoo penguins and king penguins.

9. 2. Key principals

The Government of South Georgia & the South Sandwich Islands (GSGSSI) has a risk-based response to HPAI in the Territory. This is based on observations of the outbreak since 2023 and scientific evidence of the disease's pathology based on samples taken from infected animals. This guidance aims to provide mitigation in the following key areas.

Human health -The World Health Organisation, (WHO) and Animal and Plant Health Agency (APHA) consider that the risk of human acquisition of HPAI, if biosecurity measures are followed, remains low. The UK Health Security Agency (UKHSA) have published a technical briefing [Investigation into the risk to human health of avian influenza \(influenza A H5N1\) in England: technical briefing 5 - GOV.UK](#)

Globally, a small number of cases have been recorded in humans, usually from where individuals are living or working in close proximity to large numbers of infected poultry. Effective use of PPE for those who are in close proximity to animals, or who have co-morbidities which place them at increased risk, helps to mitigate this risk. No cases of HPAI have been detected in humans on SGSSI.

Behavioural stress for wildlife – if individual animals are infected with any disease, reducing stressors is an important step to help improve chances of survival. **Reducing non-priority scientific studies that involve animal handling or approach, and ensuring responsible wildlife viewing by visitors, helps to minimise this stress factor.**

Biosecurity – the primary means of transmission of HPAI is by natural pathways but movement of infected material through human activities remains a risk. As such, maintaining separation from infected or dead animals when moving around a site and thorough cleaning of clothing, footwear, and equipment when moving between sites can help mitigate any additional risk posed by human activities.

HPAI is now widespread on South Georgia and a large amount of site- and species-specific information is available. The system of site-specific tiered restrictions that was utilised in 23/24 has been replaced with an activity-specific response measure that applies throughout the Territory. This guidance is on the basis that the following three conditions are met:

- The WHO and APHA consider that the risk to human health from the virus remains low and UKHSA advises that the protocols being implemented on South Georgia are appropriate;
- Mitigation measures in the form of specified wildlife approach distances and (for authorised science) animal handling guidelines, are in place and strictly adhered to;
- Rigorous biosecurity is in place and is enforced when arriving on South Georgia and moving around and between sites.

If at any point the assessment changes and any of these conditions are not met, this guidance will be updated, and a more restrictive approach may be adopted.

9.3 Activity specific response measures

Activity	Response measures
All	Enhanced biosecurity procedures must be followed (see 9.4)
Tourism	Prior to entering an area of high wildlife density, a trained or experienced guide or member of the research team familiar with bird behaviour should undertake a minimum 5-minute observation period of the area. For tourism visits, this should be part of the recommended 30-minute general site observation which is made prior to each landing. In this time, they should note any signs of HPAI including individuals displaying HPAI symptoms (see 9.5) and areas of high mortality*, so they can manage their visit accordingly.

	* high mortality would include areas where the density of carcasses was such that it was not possible to maintain more than 5m separation
	When ashore, staff and visitors should stay more than 5m from all live wildlife and carcasses . If it is not possible to maintain this separation, the landing should be aborted
	If HPAI symptomatic individuals are noted, consideration should be given to aborting the landing or utilizing a different part of the site. To minimise behavioural stress for wildlife, it is recommended to maintain a separation of 10m from symptomatic individuals .
	After visiting an area of high wildlife density, staff and visitors are advised to follow good hand hygiene practices (alcohol hand sanitiser or wash with soap and water)
	At sites with suspected HPAI, individuals may choose to wear a face covering. Care must be taken to avoid this becoming litter.
	If any signs of HPAI are observed, a report form should be completed and returned to GSGSSI and IAATO where applicable.
	Recognising that witnessing high levels of mortality may be distressing for visitors and the need for ELs to plan their time on SG, a list of sites where landings have been aborted due to HPAI will be maintained. This will be sent to ELs on entry to the MZ.
	At Grytviken, standard HPAI mitigation measures should be followed. Government Officers will advise on arrival if there are any areas of the site which should be avoided due to HPAI concerns.
RAP – animal handling or approach <5m	Providing there are no individuals displaying HPAI symptoms in a colony/breeding site or ongoing above baseline mortality, work may take place if 1) an RAP is in place 2) appropriate PPE is worn (see 9.6) and, 3) there is ethical (AWERB) approval.
	No animal handling or approach <5m on individuals that are displaying HPAI symptoms or for which there is ongoing, above baseline mortality within the colony/breeding site.
	For species where work has paused due to HPAI, work may recommence in a phased manner with agreement of GSGSSI providing 1) no mortality for 7 days at the colony/breeding site* 2) no individuals displaying HPAI symptoms * if mortality has been observed, this should be discussed with GSGSSI to assess if above baseline
RAP – sampling for HPAI	Collection of samples from species or environments suspected to have HPAI may only be done with a valid RAP by personnel from GSGSSI, BAS or APHA or other agencies with express permission from GSGSSI. Appropriate safety risk assessment and PPE (see 9.6) must be worn. Samples / reporting to WOH should be via the UK following testing at APHA.

9.4. HPAI Biosecurity

The following standard biosecurity measures are always in place within SGSSI.

- All biological material should be removed from boots and outer clothing and, non-porous surfaces should be treated with an appropriate biocide such as Virkon S, Bioguard, or soap followed by 3.5% bleach. This should be done before arrival to SG and between every site.*
- If clothing or equipment cannot be adequately cleaned, it should not be brought to SGSSI.
- Maintain the recommended minimum of at least **5 m** separation from all wildlife and carcasses unless you have a specific Regulated Activity Permit (RAP) which allows you to approach closer.
- Do not touch any dead wildlife or make any collections unless you have a RAP which allows this.

In addition, the following **enhanced biosecurity measures** are relevant during this period of heightened HPAI risk.

- Extra caution must be taken when cleaning any clothing or equipment which has been used in other wildlife areas especially at other sites and/or regions where HPAI has been confirmed/suspected.
- **Do not sit, kneel, crouch or lie on the ground.** Equipment should not be placed on the ground unless it is unavoidable or on a designated clean area.
- Use of tripods and monopods should be avoided, and walking poles must be cleaned thoroughly between each site.
- Remain vigilant for signs of HPAI and be prepared to abort if any behavioural signs of mortality above baseline are noted.
- After visiting an area of high wildlife density, staff and visitors are advised to follow good hand hygiene practices (alcohol hand sanitiser or wash with soap and water).

* Biocides should be used according to the manufacturer's instructions, and waste disposed of responsibly.

9.5. Signs and symptoms

Prior to entering an area of high wildlife density, a trained or experienced guide or member of the research team familiar with bird and seal behaviour should undertake a minimum 5-minute observation period of the area. For tourism visits, this should be part of the recommended 30-minute general site observation which is made prior to each landing. This may be done from a small boat, an elevated viewpoint, or (only if an appropriate Regulated Activity Permit is in place and all guidelines are suitably followed) from a drone.

The following are known signs of HPAI

- Neurological issues such as seizures, loss of coordination and balance,
- Trembling head and body,
- Sudden and rapid increase in the number of birds found dead between visits,
- Swollen head,
- Closed and excessively watery eyes. Whitening of the cornea has been noted as a potential sign in fur seals,
- Lethargy and depression, unresponsiveness, lying down, drooping wings, dragging legs,
- Twisting of the head and neck,
- Haemorrhages on the shanks of the legs and under the skin of the neck,
- Respiratory distress such as gaping (mouth breathing), nasal snicking (coughing sound), sneezing, gurgling or rattling.

NOTE: In 2024/25, a significant number of fur seals who were suspected to have HPAI had corneal edema (cloudy eyes).

9.6. Personal Protective Equipment (PPE) for science activities

Biosecurity and PPE recommendations have been scaled and are proportionate with risk. They are designed both to protect individuals and prevent spread between sites and species. For science activities, minimum PPE requirements will be identified in project specific RAPs, but individuals may choose to adopt a more precautionary approach.

Level 1 PPE - Fluid resistant face mask, disposable coveralls or outerwear i.e. oilskin jacket and trousers and eye protection to be worn. Disposable gloves or alcohol hand cleaner used between individuals. Footwear and outerwear to be decontaminated between sites/areas of high wildlife density with appropriate biocide.

Level 1 PPE will be appropriate for the majority of work which involves working in close proximity (<5m) or handling wildlife

Level 2 PPE - Tyvek 400 / fluid resistant suit, FFP2 mask and visor/eye protection or Sundstrom respirator, non-porous boots. Double-layered disposable latex gloves used between individuals and the outer glove changed each time. Footwear and outerwear to be decontaminated between sites/areas of high wildlife density with appropriate biocide.

Level 2 PPE is required for some specific high-risk activities which involve handling or invasive sampling of animals suspected of HPAI. For some types of sampling e.g. necropsies involving brain tissue, additional PPE such as respirators may be required.

Individuals must don and remove PPE correctly for it to be effective. All individuals should complete training before using level 1 or level 2 PPE. The World Health

Organisation offers online training. See - <https://openwho.org/courses/IPC-PPE-EN?locale=en>

9.7. Bird strike

If bird strike occurs on a vessel or on shore, depending on the level of injury, it may not be possible to determine if the bird has signs of HPAI or not. It is also known that some birds carry HPAI but have no visible symptoms. Therefore, a precautionary approach should be adopted.

- If birds are still alive, appropriately trained staff wearing the highest level of PPE available release the bird over the side of the vessel.
- Where organisational risk assessments and SOPs are in place, and following agreed guidelines, dazed but otherwise uninjured birds with no sign of disease may be held overnight for recuperation. Appropriate PPE must be worn.
- Any materials (used PPE, boxes etc) which have come into contact with birds should be dealt with as biosecurity waste.
- If birds are dead, carcasses should be disposed of via incineration (where available), as biological waste, or disposed of overboard by staff wearing the highest level of PPE possible.
- All bird strikes should be reported to the Government Officers on the GSGSSI bird strike pro-forma.

Guano from birds which have landed on the deck should be cleaned away using a low pressure hose/bucket and the area treated with Virkon S or an appropriate biocide.

9.8. Removal of carcasses

Collection of carcasses risks disturbance at the colony and may cause the spread of the virus and is thought to be an ineffective mitigation measure. Carcasses should only be removed/moved from a site for the purposes of post-mortem analysis for confirmation of infection, or in the station footprint of KEP or Bird Island where there is a risk to human health. This should only be done by GSGSSI or BAS staff named on an appropriate RAP who are wearing level 2 PPE.

9.9 Report form for suspected disease outbreak in wild birds



Government of South Georgia & the South Sandwich Islands

Report for suspected Highly Pathogenic Avian Influenza (HPAI) in wild birds and mammals

Name of reporter	
Name of vessel (if applicable)	
Contact details	
Site name	
Site description (lat/lon, extent of affected area, habitat type)	
Date of visit	
TIME OF REPORT (UTC)	
Length of observation period	
HPAI symptoms	
Species affected	
Approximate number of animals (per species / life stage)	
Description and number of any symptomatic animals (behaviours etc)	
Unexpected mortality	
Species affected	
Approximate number of animals (per species / life stage)	
Estimation/description of time since death i.e. fresh, skeletal remains only, mixture	
Estimation of what proportion of animals were affected	
Photos / video (as attachment) (yes/no)	
Any additional relevant information (environmental conditions etc)	

Did you abort landing as you were unable to maintain separation* from ?HPAI animals	Yes	No
Did you abort landing for another reason?	Yes	No
Please give details		

* Staff and visitors must stay more than **5m** from all **live wildlife** and **carcasses**. It is recommended to maintain a separation of **10m from symptomatic individuals**

Please return to go@gov.gs

Reports should be submitted as soon as practical, ideally within 24 hours.

Annex 1 - vehicle cleaning checklist

Before submitting your vehicle for inspection prior to cargo loading, make sure you can answer YES to all these questions:

- Has the under -carriage been inspected and is it free of soil and plant material?
- Are all tyres (including the spare) clean with particular care paid to deep treads?
- Has the engine compartment been checked (and steam cleaned where possible) to ensure there are no rodents, invertebrates or signs they may have been there?
- Have all tyres, tracks/ skis been cleaned and washed with Virkon S?
- Has the interior, including upholstery, underneath seats and storage compartments been cleaned and any biological material removed?
- Has the vehicle been fumigated with a pyrethrum based insecticide?

Vehicle description	
Name of consignee	
Organisation	
Date	

I certify that I have cleaned the above vehicle(s) as described in the GSGSSI biosecurity handbook and to the best of my knowledge, it is free from all biological material.

I understand that failure to comply with biosecurity protocols may result in a delay in my vehicle being loaded.

Signed (consignee)

Date:

Internal use only

<i>Vehicle checked on (date)</i>	
<i>Action taken</i>	
<i>Signed</i>	
<i>Date</i>	

Annex 2 Government of South Georgia & the South Sandwich Islands

Number and placement of rodent bait boxes on vessels



We have switched from traditional toxic rodent bait, to a non-toxic formulation. The non-toxic bait is highly palatable to rats and mice. It is advantageous as it avoids the risk of non-target mortality, reduces the chance that animals become 'bait-shy'/de-sensitised to toxic bait if it is needed, and reduces the amount of harmful toxins that could end up in the environment.

It is a safer and more effective method for monitoring; however **it should be noted that non-toxic bait will only indicate the presence of rodents and cannot be used to deal with a rodent problem.** The aim of monitoring is not to eliminate a rodent infestation. Monitoring tools are only there to tell us if there is an issue – if a vessel has signs of rodents on board, further additional measures will be agreed with GSGSSI to manage the problem.

Minimising the risk of rodents getting on board is essential, and includes the use of well-fitting rodent guards on mooring lines at gateway ports prior to arrival in SGSSI.

The new non-toxic bait is yellow and has an ingredient which fluoresces under ultraviolet light; not only does this better show crumbs of bait, but also stains rodent faeces and urine so they may be more easily detected.




What to Do

You will be provided with a copy of this document along with the pre-baited boxes or stations and a key to open them with. The boxes should be deployed as soon as possible (according to the instructions), and at least 24 hours prior to entering the SGSSI Maritime Zone. The boxes should remain on board throughout your visit to South Georgia, and for vessels which will return to South Georgia again, should remain in place until the next visit.

On entry to the Maritime Zone, you will be asked to declare if your vessel is rodent-free. To do that, you must first check all the bait stations and look to see if the bait has been nibbled or damaged; If it has, you must report it. The yellow non-toxic bait is perfectly safe to handle. You should also check to see if there are any other signs of rodents on board such as chewed food packaging or droppings in the galley or waste storage area. On arrival at Cumberland Bay, Government Officers will check the bait stations again.

What to Look For

Check the bait block for any signs of chew marks; it should have fairly crisp edges; if it has become rounded, something may have chewed it. Rodents may leave paired grooves or tooth marks in the surface of the bait. Look for other rodent signs nearby, such as faeces and chewed food, packaging and wood.



The picture (left) shows the relative size and shape of rat and mouse faeces, (right) shows a non-toxic block that has been nibbled by mice. Rats will leave heavier chew marks and take more (or all) of the bait.

A number of discrete areas on vessels are more favourable to rodents for entry, exit, or to live should they become established on board. These areas should be the focus of monitoring.

Rodent monitoring stations should be placed in quiet, sheltered areas, and fixed where necessary to prevent movement in heavy seas. Stations should not be placed on deck where they will get drenched by breaking seas.

Table 1 Locations of Rodent Monitoring Stations (for vessels LOA >24 metres)

Location	Type
1 Foc'sle (mooring line locker or Bosun's locker)	Rodent entry / exit point
2 Aft mooring deck	Rodent entry / exit point
3 Zodiac storage or shelter deck	Rodent entry / exit point
4 Cargo receiving areas	Rodent entry / exit point
5 Dry food & provision stores	Rodent harbourage
6 Waste storage areas	Rodent harbourage

Vessels with a load line length of more than 24 metres:

Rodent monitoring stations should be placed in each of the areas described in Table 1 where those areas exist, such that a vessel may have up to a maximum of 6 rodent monitoring stations. Note that some vessels will not have all of these distinct areas, and will require fewer bait boxes.

Yachts less than 24 metres LOA:

Yachts should place 1 rodent monitoring station inside the vessel in an area with suitable conditions for rodents, and 1 rodent monitoring station on a sheltered area of the deck while at anchor or alongside.

Rodent monitoring stations can be collected from the GSGSSI offices in Stanley, Falkland Islands, prior to departure for South Georgia.

If the vessel will not be calling in the Falklands prior to landing in South Georgia, it is the responsibility of the vessel to ensure that alternative rodent monitoring stations are in place, and that these are acceptable to GSGSSI.

If you need to source your own bait boxes and bait, you must use solid bait blocks, which are non-toxic, for example, [Detex with Lumitrack](#). Toxic solid wax bait blocks are only acceptable where you cannot procure alternative non-toxic bait. This will need to be exchanged upon the first visit to Grytviken or Stanley. Loose bait such as pellets or grains are unsuitable as they will not readily show signs of disturbance by rodents. You can also elect to create your own, such as [these](#).

GSGSSI Warehouse Biosecurity SOP

Introduction

The warehouse will be used by BAS and GSGSSI to store a range of items and for cargo processing to and from South Georgia and the South Sandwich Islands (SGSSI). It may also be used for BAS cargo storage and processing for other BAS stations in the Antarctic. It is imperative to ensure the warehouse is kept clear of biosecurity risks including any rodents, invertebrates, seeds, soil/biological material.

This document details the actions that will be taken to prevent biosecurity risks in the warehouse and what monitoring is in place to pick up breaches. A plan to deal with incursions of rodents or invertebrates into the warehouse is also described below.

This document does not include the handling of cargo, which has a separate SOP.

Prevention and Monitoring:

To prevent biosecurity risks, the following actions will be implemented:

- All cargo will be kept away from the walls (ca. 600 mm) to allow clear access around the edges for the rodent detector dog team and monitoring checks on boxes;
- Biosecure (in line with the guidance in the biosecurity handbook) cardboard boxes ready for shipping to SG should be stored off the ground on plastic pallets
- No food will be stored in the warehouse unless sealed within rodent-proof boxes (plastic);
- No food to be eaten in the warehouse;
- All rubbish should be removed from the warehouse.
- Only biosecured (biosecured in line with the guidance in the biosecurity handbook) cargo should be stored in the out-going cargo area to reduce risks of cross-contamination.

Rodent monitoring:

- Six rodent snap-trap boxes should be placed indoors around the perimeter of the warehouse (see figure 1). These traps should be checked every two weeks and snap traps inside (for wooden ones this is two snap traps and for a plastic tunnel this is one). Bait to be exchanged monthly. Bait should be a known rodent attractant, this is currently peanut butter.
- Five rodent bait boxes are placed outdoors around the perimeter of the warehouse (see figure 1). Bait to be checked monthly. Bait must be non-toxic and exchange annually or if wet/degraded. If rodents are detected inside the warehouse, the non-toxic bait should be exchanged for toxic bait (see below).
- Rodent dog team will search interior at least every month;

Invertebrates:

- Six sticky insect pads will be placed in the warehouse. One near the cargo storage areas and towards the back of the warehouse for invertebrate monitoring. These should be monitored monthly and exchanged as and when needed, a minimum every 2 months.

- Inside perimeter of outbound cargo storage area to be sprayed with residual insecticide, such as Dethlac or other broad-spectrum insecticide containing Pyrethrins and/or deltamethrin.
- External earwig spray twice a year.

Weeds:

- Areas around the building to be kept clear of weeds. Weed control is contracted once a year.
- Where possible, warehouse doors should be kept closed or only open a small amount.
- Warehouse floor sweeps to catch any seeds/windblown debris.




Figure 1: Warehouse Layout. Green squares are outdoor rodent monitoring boxes. Yellow squares are Indoor rodent monitoring boxes and insect monitoring traps.

Record Keeping:

The following spreadsheet available: <INTERNAL LINK>

It should be maintained and updated as necessary to show when tasks have been completed.

Records must be kept on:

Action	Frequency	Action Required by
Indoor bait box checks	every two weeks and to tie in 1 week ahead of Pharos sailing	Biosecurity Dog Programme
Outdoor bait box checks	TBD	Operations Manager
Bait box (indoor and outdoor) bait changes	Monthly	Biosecurity Dog Programme
Biosecurity Dog checks	monthly and to tie in 1 week ahead of Pharos sailing	Biosecurity Dog Programme

Indoor Insect Trap Checks	monthly to tie in 1 week ahead of Pharos sailing	Biosecurity Programme Dog
Insect trap glue updates	Quarterly unless there is an incursion of high-risk insects	Operations Manager/Biosecurity Dog Programme
Earwig Spraying	November and March	Operations Manager
Annual Weed Spray		Operations Manager
Sweeping	Every month and at least within 1 week after a Pharos sailing.	Operations Manager

Incursion Response

Rodents

Rodent activity around the warehouse is known and documented.

If heightened rodent activity is detected either by the presence of a mouse nest, a visible rodent in the warehouse or an indication by the detection dog team inside the warehouse.

The following actions will be taken:

- All efforts will be made to confirm the scale of incursion as appropriate (photos/dog check/additional monitoring such as fluorescent gel, camera traps etc);
- Toxic bait boxes will be deployed;
- All outbound cargo to be biosecured in line with the guidance for cargo in the biosecurity handbook and placed in a container for quarantine;
- Cargo to be searched by dog team;
- Container to be sealed with a rodent bait box containing non-toxic bait inside.
- Checks of rodent bait boxes to be checked every two days for a minimum of 14 days.
- After 14 days, continue checking until there is no sign of rodents for three checks in a row;
- Once 14 days or 3 clear checks, toxic bait be removed and snap traps to continue to be monitored weekly for 3 weeks/one additional Pharos rotation;
- Provided no further heightened rodent activity, can then move back to normal SOP.

Invertebrates:

The following is considered an invertebrate incursion:

Earwigs, spiders, ants etc – more than 3

Cockroach – more than 1

Ladybirds – more than 1

Moths - more than 1

Beetles – more than 3

Springtails – more than 1

If sticky trap checks show any of the above thresholds are breached, the following actions will be taken:

- Hand searches to take place both around the perimeter of the warehouse and on all cargo.
- Ensure all cargo is sealed – no new cargo to be placed in outbound area until the response is closed out.
- Warehouse to be fumigated with a broad-spectrum insect fogger including cyphenothrin and/or imiprothin or similar compounds such as Insecto, and the perimeter sprayed with insecticides such as Dethlac described above and left sealed for 24 hours.
- Four new traps placed inside warehouse and given 72 hours to detect any remaining live adult invertebrates.
- Traps checked and if no further incursion found, step down to routine monitoring. ~~Close out.~~
- If further incursions are found. All warehouse storage to be searched for potential and treated. Outbound cargo removed from warehouse and stored in container with insect monitoring trap. Warehouse then to be fumigated again and sealed for 24 hours.
- Heightened monitoring of traps weekly to be maintained for 3 weeks. Then proceed to normal SOP.